

dS/CFT Correspondence in Two Dimensions¹

Scott Ness² and George Siopsis³

Department of Physics and Astronomy,
The University of Tennessee, Knoxville
TN 37996 - 1200, USA.

Abstract

We discuss the quantization of a scalar particle moving in two-dimensional de Sitter space. We construct the conformal quantum mechanical model on the asymptotic boundary of de Sitter space in the infinite past. We obtain explicit expressions for the generators of the conformal group and calculate the eigenvalues of the Hamiltonian. We also show that two-point correlators are in agreement with the Green function one obtains from the wave equation in the bulk de Sitter space.

¹Research supported in part by the DoE under grant DE-FG05-91ER40627.

²ness@utk.edu

³gsiopsis@utk.edu

Even though the AdS/CFT correspondence is by now well-understood [1–8], a similar correspondence for a de Sitter space has been quite a puzzle to establish. Such a development is of considerable interest in view of recent astronomical data suggesting that we live in a Universe of positive cosmological constant [9].

A concrete proposal for a dS/CFT correspondence was recently put forth by Strominger [10] and attracted much attention [11]. According to this proposal, all observables in the bulk de Sitter space are generated by data specified on its asymptotic boundary which can be selected to be the Euclidean hypersurface $\mathcal{I}^-$ in the infinite past. The isometries of the de Sitter space are mapped onto generators of the conformal group of the theory defined on the boundary $\mathcal{I}^-$. This CFT is hard to construct in general, but various features, such as conformal weights and masses, are known. In three-dimensional de Sitter space, the conformal group on the boundary is infinite and the central charge is known [12]. The CFT is a Liouville theory [13].

Here we discuss the case of two-dimensional de Sitter space. The asymptotic boundary $\mathcal{I}^-$ is a circle, which upon a Wick rotation turns into time. The theory on the boundary is a conformal quantum mechanical model. We shall explicitly construct this model for the case of a scalar particle, obtain the generators of the conformal group, calculate the eigenvalues of the Hamiltonian and the Green functions. Our method of solution is similar to the one discussed by de Alfaro, Fubini and Furlan (DFF) [14], even though our Hamiltonian differs from theirs.

The two-dimensional de Sitter space (dS_2) may be parametrized as

$$\frac{ds^2}{l^2} = -d\tau^2 + \cosh^2 \tau d\phi^2. \quad (1)$$

We will explicitly consider the case where ϕ spans the real axis (zero temperature limit) and then comment on what changes need to be made to turn ϕ periodic (finite temperature).

Consider a scalar field Φ of mass m . It obeys the wave equation in de Sitter space

$$l^2 \nabla^2 \Phi = -\frac{1}{\cosh \tau} \partial_\tau (\cosh \tau \partial_\tau \Phi) + \frac{1}{\cosh^2 \tau} \partial_\phi^2 \Phi = m^2 l^2 \Phi. \quad (2)$$

The operator ∇^2 is the Casimir operator in the $SL(2, \mathbb{R})$ algebra generated by

$$L_\pm = e^{\pm i\phi} \left(i \tanh \tau \frac{\partial}{\partial \phi} \pm \frac{\partial}{\partial \tau} \right), \quad L_3 = -i \frac{\partial}{\partial \phi} \quad (3)$$

Indeed,

$$\ell^2 \nabla^2 = \frac{1}{2} (L_+ L_- + L_- L_+) - L_3^2 \quad (4)$$

Focusing on the $\mathcal{I}^-$ boundary reduces the wave equation to

$$(-\partial_\tau^2 + \partial_\tau + 4e^{2\tau}\partial_\phi^2)\Phi = m^2 l^2 \Phi. \quad (5)$$

On this spacelike slice we see the third term is negligible and the wave equation becomes independent of ϕ . The solution to the wave equation (2) behaves asymptotically as

$$\Phi \sim e^{h_\pm \tau}, \quad h_\pm = \frac{1}{2} \pm \nu, \quad \nu = i\mu, \quad \mu = \frac{1}{2}\sqrt{4m^2 l^2 - 1}. \quad (6)$$

We will concentrate on the case of imaginary ν ($\mu \in \mathbb{R}$) in which $h_- = h_+^*$. Introducing the coordinate $q = e^\tau$, we may write (5) as

$$f_k'' + 4k^2 f_k = -\frac{m^2 \ell^2}{q^2} f_k \quad (7)$$

where $\Phi(\tau, \phi) = e^{ik\phi} f_k(q)$. The solutions are

$$f_k(q) = \sqrt{2kq} Z_\nu(2kq) \quad (8)$$

where Z_ν is a Bessel function. They form an orthogonal set under the inner product

$$(\Phi_1, \Phi_2) = i \int_{-\infty}^{\infty} \frac{d\phi}{2\pi} e^{i(k_1 - k_2)\phi} (f_{k_1}^*(q) f_{k_2}'(q) - f_{k_1}'^*(q) f_{k_2}(q)). \quad (9)$$

The apparent q -dependence disappears after we apply the wave equation and the integral over ϕ leads to a δ -function, as expected. For the Euclidean choice

$$\Phi_k^E = C^E e^{ik\phi} \sqrt{2kq} H_\nu^{(1)}(2kq) \quad (10)$$

demanding orthonormality, fixes the normalization constant to $C^E = \frac{1}{\sqrt{8}}$ for real ν , where we used the Wronskian $H_\nu^{(1)}(x)H_\nu^{(2)'}(x) - H_\nu^{(1)'}(x)H_\nu^{(2)}(x) = \frac{-4i}{\pi x}$. Another interesting choice is

$$\Phi_k^+ = C^+ e^{ik\phi} \sqrt{2kq} J_\nu(2kq). \quad (11)$$

For an orthonormal set, we choose

$$C^+ = \frac{\sqrt{\nu}}{2} \frac{\Gamma(\nu)}{\Gamma(\frac{1}{2})} \quad (12)$$

Using the Bessel function identity

$$J_\nu(x) = \frac{1}{2} (H_\nu^{(1)}(x) - e^{i\pi\nu} H_\nu^{(1)}(-x)) \quad (13)$$

we obtain

$$\Phi_k^+(q, \phi) = \sqrt{\frac{\nu}{2\pi}} \Gamma(\nu) (\Phi_k^E(q, \phi) + e^{i\pi h_+} \Phi_k^E(-q, \phi)) \quad (14)$$

which differs from [12] by a phase. The Green function for the modes Φ_k^+ can be obtained from

$$G^+(q, \phi; q', \phi') = \int \frac{dk}{k} \Phi_k^+(q, -\phi) \Phi_k^+(q', \phi'). \quad (15)$$

After some algebra, we arrive at

$$G^+(q, \phi; q', \phi') = (C^+)^2 \frac{\Gamma(h_+)}{\Gamma(h_+ + \frac{1}{2})\Gamma(\frac{1}{2})} (4P)^{-h_+} F(h_+, h_+; 2h_+; -1/P) \quad (16)$$

where P measures the distance between the two arguments of the Green function in the three-dimensional Minkowski space in which dS_2 is embedded,

$$P = \frac{4(q - q')^2 + (\phi - \phi')^2}{16qq'}. \quad (17)$$

When $P \rightarrow \infty$, the Green function behaves as

$$G^+(q, \phi; q', \phi') \sim (C^+)^2 \frac{\Gamma(h_+)}{\Gamma(h_+ + \frac{1}{2})\Gamma(\frac{1}{2})} (4P)^{-h_+}. \quad (18)$$

For the Euclidean modes, the Green function can be found in terms of the Green function obtained above using (14) [10]. Near the boundary, we deduce the two-point function of the dual conformal theory,

$$\langle \mathcal{O}^\dagger(\phi) \mathcal{O}(\phi') \rangle = 2\sqrt{\pi} \frac{\Gamma(h_+)}{\Gamma(\nu)} (\phi - \phi')^{-2h_+} \quad (19)$$

If ϕ is a periodic coordinate with period 2π , then k takes on discrete values and the integral defining the Green function turns into a sum. The form of the two-point function may be deduced from the periodicity condition and the singularity, which uniquely determine it. We obtain

$$\langle \mathcal{O}^\dagger(\phi) \mathcal{O}(\phi') \rangle = 2\sqrt{\pi} \frac{\Gamma(h_+)}{\Gamma(\nu)} \left(2 \sin \frac{\phi - \phi'}{2} \right)^{-2h_+} \quad (20)$$

The boundary conformal theory can be defined in terms of the coordinates $q = e^\tau$, ϕ and their conjugate momenta $p = e^{-\tau} \partial_\tau$, H , respectively. Instead of the $SL(2, \mathbb{R})$ generators (3), it is more convenient to introduce the generators

$$\begin{aligned} D &= i(\phi \partial_\phi + \partial_\tau) \\ K &= -\phi^2 \partial_\phi + 2i\phi \partial_\tau + 4e^{2\tau} \partial_\phi. \end{aligned} \quad (21)$$

in the vicinity of the boundary $\mathcal{I}^-$. The three operators H, D and K satisfy the $SL(2, \mathbb{R})$ algebra

$$[H, K] = 2iD, \quad [H, D] = iH, \quad [K, D] = -iK. \quad (22)$$

The wave equation is mapped onto the constraint

$$4q^2 H^2 - q^2 p^2 = g \quad , \quad g = m^2 \ell^2 + A \quad (23)$$

where we have allowed for the possibility of a normal ordering ambiguity by introducing the constant A . Classically, we can solve the constraint (23) and express the conformal Hamiltonian as

$$H = \frac{1}{2} \sqrt{p^2 + \frac{g}{q^2}}. \quad (24)$$

The constraint thus reduces the system to a quantum mechanical model with a single degree of freedom q , with ϕ playing the role of imaginary time. Notice that our Hamiltonian differs from the one considered by DFF [14] (theirs is the square of ours), yet the results are similar.

In order to quantize this system, we promote (q, p) to operators (Q, P) and impose the standard commutation relation

$$[Q, P] = i \quad (25)$$

The coordinate ϕ is identified with time after a Wick rotation. D and K (eq. (21)) become the dilation and special conformal operators, respectively, given by

$$\begin{aligned} D &= i \left(\phi H + \frac{1}{2} (QP + PQ) \right) \\ K &= -\phi^2 H + i\phi(QP + PQ) + 2(Q^2 H + H Q^2). \end{aligned} \quad (26)$$

Following the procedure in [14], it is more convenient to start with the values of the $SL(2, \mathbb{R})$ generators at $\phi = 0$. Let $Q_0 = Q(0)$ and $P_0 = P(0)$. The three generators reduce to

$$\begin{aligned} H &= \frac{1}{2} \sqrt{P_0^2 + \frac{g}{Q_0^2}} \\ D &= \frac{i}{2} (Q_0 P_0 + P_0 Q_0) \\ K &= 2(Q_0^2 H + H Q_0^2). \end{aligned} \quad (27)$$

It can be checked explicitly that they obey the $SL(2, \mathbb{R})$ algebra

$$[H, K] = -2iD, \quad [H, D] = -iH, \quad [K, D] = iK. \quad (28)$$

The Casimir operator of this algebra is given by

$$N = \frac{1}{2}(HK + KH) - D^2. \quad (29)$$

This commutes with all operators and its numerical value should equal $m^2\ell^2$ to match the wave equation. To calculate N , we first bring it into the form

$$N = [H, [Q_0^2, H]] + g + \frac{3}{4} \quad (30)$$

after some straightforward operator manipulations. The remaining double commutator is independent of g and can therefore be easily calculated by setting $g = 0$ in H . It may also be deduced from the corresponding Poisson brackets. The final result is

$$N = g - \frac{1}{4} \quad (31)$$

By comparing with the wave equation (5), we deduce that the normal ordering constant is $A = 1/4$ (*cf.* (23)) and

$$g = m^2\ell^2 + \frac{1}{4} \quad (32)$$

We follow the procedure of [14] to construct the states in the Hilbert space. Define new operators R, S :

$$\begin{aligned} R &= \frac{1}{2} \left(\frac{1}{a} K + aH \right), \\ S &= \frac{1}{2} \left(\frac{1}{a} K - aH \right). \end{aligned} \quad (33)$$

where a is a constant with dimension of length. We will let $a = 1$ for simplicity. R plays the role of a Hamiltonian for the quantum mechanical system on $\mathcal{I}^-$ whose eigenstates span the Hilbert space. By including the dilation operator, R, S and D form the closed algebra $O_{2,1}$,

$$[D, R] = iS \quad [S, R] = -iD \quad [S, D] = -iR. \quad (34)$$

Next, we define raising and lowering operators

$$L_{\pm} = S \mp iD. \quad (35)$$

These new operators, along with R form the closed algebra $SL(2, \mathbb{R})$,

$$[R, L_{\pm}] = \pm L_{\pm} \quad [L_+, L_-] = -2R. \quad (36)$$

The Casimir operator for this algebra (29) can be written in terms of R, L_+ and L_- as

$$N = \frac{1}{2}(HK + KH) - D^2 = R^2 - R - L_+L_- \quad (37)$$

The ground state energy can be found by operating on the ground state of R ,

$$R|0\rangle = r_0|0\rangle, \quad (38)$$

$$N|0\rangle = r_0(r_0 - 1)|0\rangle = m^2 l^2 |0\rangle. \quad (39)$$

Thus, the ground state energy is

$$r_0 = \frac{1}{2}(1 \pm \sqrt{1 - 4m^2 l^2}). \quad (40)$$

Therefore, $r_0 = h_{\pm}$ (eq. (6)). Without loss of generality, we choose $r_0 = h_+$. The other choice ($r_0 = h_- = h_+^*$) leads to a dual Hilbert space related to the one we are about to study via complex conjugation. It should be noted that inner products ought to be defined in terms of both Hilbert spaces, leading to normalization conditions such as (44).

By successively acting with the raising operators, we deduce the eigenvalues of R ,

$$R\psi_n(q) = (r_0 + n)\psi_n(q) \quad (41)$$

In the position representation, we obtain the Schrödinger equation

$$\left((1 + 2q^2) \sqrt{-\frac{d^2}{dq^2} + \frac{g}{q^2}} + 2 \sqrt{-\frac{d^2}{dq^2} + \frac{g}{q^2}} q^2 \right) \psi_n(q) = \sqrt{2}(r_0 + n)\psi_n(q). \quad (42)$$

To solve the Schrödinger equation, it is convenient to introduce the transition matrix $\mathcal{T}_{nk}$ between the two bases Φ_k^+ (11) and ψ_n , at time $\phi = 0$,

$$\psi_n(q) = \int_0^\infty dk \mathcal{T}_{nk}^* \Phi_k^+(q), \quad (43)$$

where we impose orthonormality,

$$\int_0^\infty dq \psi_n(q) \psi_{n'}(q) = \delta_{nn'}. \quad (44)$$

The action of the three $SL(2, \mathbb{R})$ generators on Φ_k^+ translates to an action on the matrix elements $\mathcal{T}_{nk}$,

$$\hat{H}\mathcal{T}_{nk} = k\mathcal{T}_{nk} \quad , \quad \hat{D}\mathcal{T}_{nk} = -i \left(k \frac{d}{dk} + \frac{1}{2} \right) \mathcal{T}_{nk} \quad , \quad \hat{K}\mathcal{T}_{nk} = \left(-k \frac{d^2}{dk^2} - \frac{d}{dk} + \frac{\nu^2}{k} \right) \mathcal{T}_{nk}, \quad (45)$$

respectively. In this representation, the Schrödinger equation (42) becomes

$$\frac{1}{2}(\hat{H} + \hat{K})\mathcal{T}_{nk} = (h_+ + n)\mathcal{T}_{nk} \quad (46)$$

which can be solved. The solution is given in terms of Laguerre polynomials [14],

$$\mathcal{T}_{nk} = \frac{2^{h_+}}{C^+} \sqrt{\frac{n!}{\Gamma(n + 2h_+)}} k^\nu e^{-k} L_n^{2\nu}(2k) \quad (47)$$

where the normalization has been fixed as in [14] with the exception of the factor C^+ which is due to the different normalization condition we imposed on the wavefunctions Φ_k^+ . It leads to a unit norm for the wavefunction ψ_n . It is convenient to introduce the Laplace transform of $\mathcal{T}_{nk}$ with respect to the energy k ,

$$\tilde{\mathcal{T}}_n(\phi) = 2^{-h_+} \int_0^\infty dk e^{-k\phi} k^\nu \mathcal{T}_{nk} \quad (48)$$

given in terms of the conjugate time variable ϕ . The additional energy factor k^ν ensures that the action of the $SL(2, \mathbb{R})$ generators is as expected. Explicitly,

$$\tilde{\mathcal{T}}_n(\phi) = \frac{(-)^n}{(C^+)^2} \sqrt{\frac{\Gamma(n + 2h_+)}{n!}} \left(\frac{1 - \phi}{1 + \phi} \right)^{n+h_+} (1 - \phi^2)^{-h_+} \quad (49)$$

and the $SL(2, \mathbb{R})$ generators act as (*cf.* eq. (45))

$$\hat{H}\tilde{\mathcal{T}}_n = -\frac{d}{d\phi}\tilde{\mathcal{T}}_n \quad , \quad \hat{D}\tilde{\mathcal{T}}_n = i \left(\phi \frac{d}{d\phi} + h_+ \right) \tilde{\mathcal{T}}_n \quad , \quad \hat{K}\tilde{\mathcal{T}}_n = \left(\phi^2 \frac{d}{d\phi} + 2h_+\phi \right) \tilde{\mathcal{T}}_n. \quad (50)$$

The two-point function is given by

$$G(\phi, \phi') = \sum_{n=0}^{\infty} \tilde{\mathcal{T}}_n(\phi) \tilde{\mathcal{T}}_n(-\phi') \quad (51)$$

Notice that we there is no complex conjugation on $\tilde{\mathcal{T}}_n(-\phi')$, because it ought to be the complex conjugate of the dual of $\tilde{\mathcal{T}}_n$ which is itself related to $\tilde{\mathcal{T}}_n$ by complex conjugation (*cf.* also the inner product (44)). A short calculation reveals

$$G(\phi, \phi') = \frac{\Gamma(2h_+)}{2^{2h_+}(C^+)^2} (\phi - \phi')^{-2h_+} = 2\sqrt{\pi} \frac{\Gamma(h_+)}{\Gamma(\nu)} (\phi - \phi')^{-2h_+} \quad (52)$$

in agreement with the de Sitter space result (19). Higher-order correlators may also be obtained by using the standard methods developed in [14].

If we demand periodicity of ϕ ($\phi \equiv \phi + 2\pi$), the above calculations become cumbersome. The energy k takes on discrete values and the differential equations determining the transition matrix $\mathcal{T}_{nk}$ turn into difference equations. Nevertheless, correlators may be deduced without explicit calculations, because they are uniquely determined by the periodicity requirements and their singularities.

To summarize, we have discussed the conformal quantum mechanical model which resides on the boundary $\mathcal{I}^-$ of two-dimensional de Sitter space in the infinite past. We calculated the eigenvalues and corresponding eigenfunctions of the Hamiltonian and deduced correlation functions. We showed that the Green functions agree with the propagators one obtains from the de Sitter space wave equation. Most of our results are similar to those of ref. [14], even though our Hamiltonian is different and the Schrödinger equation (42) cannot be solved explicitly. What saves the day is the $SL(2, \mathbb{R})$ symmetry which determines much of the structure of the conformal theory as was shown in [14]. It would be interesting to extend our results to de Sitter spaces of dimension higher than two and shed some light on the behavior of the conformal field theories on the boundary.

References

- [1] J. D. Brown, M. Henneaux, “Central Charges In the Canonical Realization Of Asymptotic Symmetries: An Example From Three-Dimensional Gravity,” *Commun. Math. Phys.* **104** (1986) 207.
- [2] V. Balasubramanian, P. Kraus, “A stress tensor for Anti-de Sitter gravity,” *Commun. Math. Phys.* **208** (1999) 413 [hep-th/9902121].
- [3] S. B. Giddings, “The boundary S-matrix and the AdS to CFT dictionary,” *Phys. Rev. Lett.* **83** (1999) 2707 [hep-th/9903048].
- [4] S. S. Gubser, “Non-conformal examples of AdS/CFT,” *Class. Quant. Grav.* **17** (2000) 1081.
- [5] E. Witten, “Anti-de Sitter space and holography”, *Adv. Theor. Math. Phys.* **2** (1998) 253 [hep-th/9802150].
- [6] G.T. Horowitz, R. C. Myers, “The AdS/CFT Correspondence and a New Positive Energy Conjecture for General Relativity,” *Phys. Rev.* **D59** (1999) 026005 [hep-th/9808079].
- [7] L. Susskind, E. Witten, “The Holographic Bound in Anti-de Sitter Space,” hep-th/9805114.
- [8] H. Boschi-Filho, N. R. F. Braga, “Compact AdS space, Brane geometry and the AdS/CFT correspondence,” hep-th/0112196.
- [9] S. Permuter, “Supernovae, dark energy and the accelerating universe: The status of the cosmological parameters,” in *Proc of the 19th Intl. Symp. on photon and lepton interactions at High Energy LP99* eds. J. A. Jaros and M. E. Peskin, *Int. J. Mod. Phys.* **A15 S1** 2000 715 [eConfC **990809** 2000 715].
- [10] A. Strominger, “The dS/CFT correspondence,” *JHEP* **0110** (2001) 034 [hep-th/0106113].
- [11] A. Strominger, “Inflation and the dS/CFT Correspondence,” hep-th/0110087;
A. C. Petkou, G. Siopsis, “dS/CFT correspondence on a brane,” hep-th/0111085.
S. Nojiri, S. D. Odintsov, “Wilson loop and dS/CFT correspondence,” hep-th/0112152.

- A. J. M. Medved, “A Holographic Interpretation of Asymptotically de Sitter Spacetimes,” hep-th/0112226.
- A. M. Ghezelbash, D. Ida, R. B. Mann, T. Shiromizu, “Slicing and Brane dependence of the (A)dS/CFT Correspondence,” hep-th/0201004.
- [12] R. Bousso, A. Maloney, A. Strominger, “Conformal vacua and Entropy in de Sitter Space,” hep-th/0112218.
- [13] S. Cacciatori, D. Klemm, “The Asymptotic Dynamics of de Sitter Gravity in three Dimensions,” Class. Quant. Grav. **19** (2002) 579 [hep-th/0110031].
- [14] V. De Alfaro, S. Fubini, G. Furlan, “Conformal Invariance in Quantum Mechanics”, Il Nuovo Cimento **34A** (1976) 569.